

# Incidence and determinants of antiretroviral switching away from TDF-based backbone in the recent years in the Iona Foundation Cohort

Vergori A<sup>1</sup>, Lorenzini P<sup>1</sup>, Cozzi-Lepri A<sup>2</sup>, Maggiolo F<sup>3</sup>, Lapadula G<sup>4</sup>, De Luca A<sup>5</sup>, Cingolani A<sup>6</sup>, Galli M<sup>7</sup>, Mazzarello G<sup>8</sup>, Milini P<sup>9</sup>, d'Arminio Monforte A<sup>10</sup>, Antinori A<sup>1</sup> on behalf of ICONA Foundation Study Group.



P100

<sup>1</sup>HIV/AIDS Unit, INMI L. Spallanzani IRCCS, Rome, Italy; <sup>2</sup>Institute for Global Health, UCL London, UK; <sup>3</sup>Division of Infectious Diseases, ASST Papa Giovanni XXIII, Bergamo, Italy; <sup>4</sup>Division of Infectious Diseases, ASST Monza University of Milano-Bicocca, Italy; <sup>5</sup>University Division of Infectious Diseases, Siena University Hospital, University of Siena, Italy; <sup>6</sup>Infectious Diseases Division, Pol. A. Gemelli, Catholic University, Rome, Italy; <sup>7</sup>3rd Division of Infectious Diseases, ASST FB-Sacco University of Milan, Italy; <sup>8</sup>Infectious Disease Clinic, IRCCS University Hospital, San Martino-IST, Genoa, Italy; <sup>9</sup>Infectious Diseases Division, General Hospital, Macerata, Italy; <sup>10</sup>Clinic of Infectious and Tropical Diseases, ASST Santi Paolo e Carlo Hospital, University of Milan, Italy

## BACKGROUND

In the past years, NRTI backbone based on tenofovir disoproxil fumarate (TDF) and emtricitabine (FTC) has represented the standard of care in the antiretroviral setting.

Increasing concerns about TDF renal toxicity, as itself or in combination with other antiretrovirals, and the availability of effective and safe alternatives, as abacavir (ABC), tenofovir alafenamide (TAF) or NRTI-sparing less-drug regimens (LDRs), enhances chances of TDF substitution.

Rate and predictors of TDF discontinuation may have public health implications in the view of availability of TDF/FTC generic formulation.

## AIM

To report data from real life on the incidence and factors associated with discontinuation of TDF in the INSTI antiretroviral era.

## STUDY DESIGN AND METHODS

HIV-1 positive patients from the Iona Foundation Cohort, aged 18 years or over, initiating their first cART regimen with TDF-based backbone plus a 3<sup>rd</sup> drug from January 2009 onwards were included. Patients were included if they had been treated for >30 days with TDF, subject with HbsAg positivity were not included.

Primary endpoint of the analysis was TDF discontinuation for any reasons. The adjusted risk of TDF discontinuation was estimated by Cox regression according to main fixed covariates at baseline: gender, race, CDC C stage, mode of HIV transmission, HCVAb status, CD4 and CD8 count, HIVRNA, total and HDL cholesterol, total number of non-communicable comorbidities (NCC) (of whom CKD, hypertension, diabetes, previous cardiovascular or hepatic event) and third drug (the class or the single drug). eGFR during TDF exposure was estimated by CKD-EPI formula and was used in the models as time dependent covariate. Calendar year of cART initiation was divided into three periods: 2009-2011, 2012-2014, 2015-2017.

## RESULTS I

5544 ART-naive patients were included in the analyses: 81% males, median age of 39 years (IQR 31-47), 10% were on a CDC stage C, median CD4 count 332 cells/mm<sup>3</sup> (178-474), median HIV RNA 4.74 (4.16-5.24) Log<sub>10</sub> cps/mL. 2296 (41.4%) started a NNRTI-based, 2015 (36.2%) a PI/b-based and 1233 (22.4%) a INSTI-based regimen. 2546 (46%) pts discontinued TDF after a median of 2.3 years (IQR 1.1-3.9).

## Acknowledgments

### ICONA Foundation Study Group

**BOARD OF DIRECTORS:** A d'Arminio Monforte (President), A Antinori (Vice-President), M Andreoni, A Castagna, F Castelli, R Cauda, G Di Perri, M Galli, R Iardino, G Ippolito, A Lazzarin, GC Marchetti, G Rezza, F von Schloesser, P Viale.

**SCIENTIFIC SECRETARY:** A d'Arminio Monforte, A Antinori, A Castagna, F Ceccherini-Silberstein, A Cozzi-Lepri, E Girardi, S Lo Caputo, C Mussini, M Puoti, CF Perno.

**STEERING COMMITTEE:** A Antinori, F Bai, C Balotta, A Bandera, S Bonora, M Borderi, A Calcagno, A Capetti, MR Capobianchi, A Castagna, F Ceccherini-Silberstein, S Cicalini, A Cingolani, P Cinque, A Cozzi-Lepri, A d'Arminio Monforte, A De Luca, A Di Biagio, E Girardi, N Gianotti, A Gori, G Guaraldi, G Lapadula, M Lichtner, S Lo Caputo, G Madeddu, F Maggiolo, G Marchetti, L Monno, C Mussini, S Nozza, CF Perno, C Pinnetti, M Puoti, E Quiros Roldan, R Rossotti, S Rusconi, MM Santoro, A Saracino, L Sarmati.

**STATISTICAL AND MONITORING TEAM:** A Cozzi-Lepri, I Fanti, L Galli, P Lorenzini, A Rodano, M Macchia, A Tavelli.

**BIOLOGICAL BANK INMI:** F Carletti, S Carrara, A Di Caro, S Graziano, F Petroni, G Proto, S Truffa.

**PARTICIPATING PHYSICIANS AND CENTERS:** Italy A Giacometti, A Costantini, V Barocci (Ancona); G Angarano, L Monno, C Fabrizio (Bari); F Maggiolo, C Suardi (Bergamo); P Viale, V Donati, G Verucchi (Bologna); F Castelnuovo, C Minardi, E Quiros Roldan (Brescia); B Menzaghi, C Abeli (Busto Arsizio); B Cocopardo, B Celestia (Catania); J Vecchiet, K Falasca (Chieti); A Pan, S Lorenzotti (Cremona); L Sighinolfi, D Segala (Ferrara); P Blanc, F Vichi (Firenze); G Cassola, C Viscoli, A Alessandrini, N Bobbio, G Mazzarello (Genova); M Lichtner, S Vita, (Latina); P Bonfanti, C Molteni (Lecco); A Chioldera, P Milini (Macerata); G Nunnari, G Pellicano (Messina); A d'Arminio Monforte, M Galli, A Lazzarin, G Rizzardini, M Puoti, A Castagna, S Cannizzo, MC Moioli, R Piolini, AL Ridolfo, S Salpietro, C Tincati, (Milano); C Mussini, C Puzzolante (Modena); C Migliorino, G Lapadula (Monza); V Sangiovanni, G Borgia, V Esposito, F Di Martino, I Gentile, L Maddaloni (Napoli); AM Cattelan, S Marinello (Padova); A Cascio, C Colomba (Palermo); F Baldelli, E Schiaroli (Perugia); G Parruti, F Sozio (Pescara); G Magnani, MA Ursitti (Reggio Emilia); M Andreoni, A Antinori, R Cauda, A Cristaudo, V Vullo, R Acinapura, G Baldin, M Capozzi, A Mondini, A Cingolani, M Rivano Capparucia, G Iaini, A Latini, R Gagliardini, MM Plazzi, S Savinelli, A Vergori (Roma); M Cecchetto, F Viviani (Rovigo); G Madeddu, P Bagella (Sassari); A De Luca, B Rossetti (Siena); A Franco, R Fontana Del Vecchio (Siracusa); D Francisci, C Di Giulii (Terni); P Caramello, G Di Perri, S Bonora, GC Orofino, M Sciandra (Torino); M Bassetti, A Londero (Udine); G Pellizzer, V Manfrin (Vicenza); G Starnini, A Ialungo (Viterbo).

## RESULTS II

The Incidence Rate (IR) of TDF discontinuation for any reason significantly increased from 10.3 (95%CI 9.5-11.1) per 100 PYFU in 2009-11, to 14.3 (13.4-15.2) in 2012-14 and to 34.9 (32.7-37.7) in 2015-17 (p<0.001). Two separate models [a) and b)] to estimate hazard risk by third drug class and by type of third drug were performed (Table 1). Using NNRTI as reference, an increased risk of TDF discontinuation was found both for PI/b (aHR 1.55; 95%CI 1.41-1.71) and INSTI (2.09; 1.85-2.35). DTG (2.78; 2.32-3.33), EVG/c (3.59; 2.94-4.39) and RAL (2.19; 1.79-2.69) also were associated with higher risk of discontinuation respect to EFV.

By multivariable Cox regression, a lower current eGFR was associated to higher risk of discontinuing TDF in all time periods. HR for eGFR<60 became considerably lower in last 3-year period (Table 2). There was a significant interaction between calendar year of cART initiation and current eGFR level (p<0.001) (Figure 1).

Non-communicable comorbidities (NCC) were associated in first and second periods, but not in third one. P at interaction test was significant (p<0.001) (Table 2).

**Table 1 – Crude and adjusted hazard ratio (HR) of TDF discontinuation for any causes by 3<sup>rd</sup> drug class and by single 3<sup>rd</sup> drug**

a)	HR	95%CI	p-value	aHR*	95%CI	p-value
NNRTI	1					
PI/b	1.36	1.25 1.50	<0.001	1.55	1.41 1.71	<0.001
INSTI	4.11	3.70 4.57	<0.001	2.09	1.85 2.35	<0.001

b)	HR	95%CI	p-value	aHR*	95%CI	p-value
EFV	1					
RPV	3.02	2.61 3.48	<0.001	1.56	1.33 1.83	<0.001
DRV/r or /c	2.48	2.17 2.83	<0.001	1.92	1.67 2.20	<0.001
ATV/r or /c	1.96	1.71 2.25	<0.001	1.92	1.67 2.20	<0.001
DTG	8.52	7.28 9.96	<0.001	2.78	2.32 3.33	<0.001
EVG	11.18	9.37 13.34	<0.001	3.59	2.94 4.39	<0.001
RAL	3.36	2.76 4.08	<0.001	2.19	1.79 2.69	<0.001
other	1.50	1.25 1.80	<0.001	1.59	1.31 1.92	<0.001

\*adjusted for gender, ethnicity, mode of HIV transmission, HCV coinfection, CD4 and CD8 count, HIVRNA, total and HDL cholesterol, number of NCC, calendar year of cART initiation and current eGFR

**Table 2 – Factors independently associated with TDF discontinuation according to three time periods (2009-11; 2012-14; 2015-17) of cART initiation. Significant results are reported in bold.**

	2009-2011		2012-2014		2015-2017	
	aHR*	95%CI	aHR*	95%CI	aHR*	95%CI
Current e-GFR ml/min/1,73 m <sup>2</sup> (by CKD-EPI)						
- >90 (ref.)	1.00		1.00		1.00	
- 60-90	1.65	1.41-1.96	1.45	1.26-1.67	1.11	0.96-1.28
- <60	4.96	3.45-7.14	4.29	3.18-5.77	1.56	1.05-2.31
NCC						
- 0 (ref.)	1.00		1.00		1.00	
- 1	1.65	1.20-2.26	1.23	0.94-1.60	1.01	0.75-1.35
- 2+	2.21	1.03-4.75	1.77	1.04-3.04	1.12	0.63-2.02
ARV third drug class						
- NNRTI (ref.)	1.0		1.0		1.0	
- PI/b	1.92	1.62-2.28	1.37	1.19-1.57	1.45	1.15-1.84
- INSTI <sup>§</sup>	1.17	0.73-1.89	2.12	1.70-2.64	2.43	2.04-2.89


\*adjusted for gender, ethnicity, mode of HIV transmission, HCV coinfection, CD4 and CD8 count, HIVRNA, total and HDL cholesterol

§ A sensitivity analysis, performed after excluding EVG/c, confirmed these results.

**Table 3 - Regimen started after TDF discontinuation**

Regimen started after TDF	Over 2224 pts who started a new regimen after TDF
TAF/FTC	1008 (45.3%)
ABC/3TC	601 (27.0%)
Other backbone	120 (5.4%)
NRTI-sparing LDRs	445 (20.0%)

**Figure 1 – Adjusted hazard ratios of tenofovir disoproxil fumarate (TDF) discontinuation according to calendar year of TDF initiation and current e-GFR estimated by CKD-Epi formula.**



Regimen started after TDF discontinuation are shown in Table 3.

## CONCLUSIONS

In our cohort, a significant increase of TDF discontinuation was found after 2015. Associated drugs (PI/b and INSTI) and eGFR decline mainly predicted drug change, with a lower risk of switching away from TDF at declining eGFR levels in the last period. The remarkable risk of TDF switching in people receiving INSTI, increasingly in the last three years, may suggest physicians attitudes towards co-formulated regimens more than TDF safety concerns in clinical decision.

## Funding

ICONA Foundation is supported by unrestricted grants from BMS, Gilead Sciences, Janssen, MSD and Viiv Healthcare

## References

1. EACS Guidelines, Version n.9,2017; 2. Linee Guida Italiane;8 edizione,2017; 3. Lee FJ, Plos One, 2014; 4. Horberg M, JAIDS 2010; 5. Mocroft A, AIDS 2010; 6. Ryom L, JID 2013; 7. Ryom L, AIDS 2014; 8. Ryom L, AIDS 2017; 9. Scherzer R, AIDS 2012; 10. Mocroft A, AIDS 2016; 11. Costarelli S, Plos One 2016; 12. Mills A, Lancet Infect Dis, 2016; 13. Trottier B, Antivir Ther, 2017; 14. Orkin C, Lancet HIV, 2017; 15. Perez-Molina JA, Lancet Infect Dis, 2015; 16. Di Giambenedetto S, JAC, 2017; 17. Pulido F, Clin Infect Dis, 2017; 18. Maggiolo F, J Acquir Immune Defic Syndr, 2016; 19. Taiwo BO, Clin Infect Dis, 2017; 20. Libbre JM, Lancet, 2018

## Contact Information

andrea.antinori@inmi.it; alessandra.vergori@inmi.it; patrizia.lorenzini@inmi.it